

AUSTRALIAN PRODUCT INFORMATION
Cipla Hydrocortisone
(hydrocortisone sodium succinate) powder for injection

1. NAME OF THE MEDICINE

Hydrocortisone sodium succinate

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 133.7 mg of hydrocortisone sodium succinate equivalent to 100 mg of hydrocortisone.

For the full list of excipients, see section 6.1 List of excipients.

3. PHARMACEUTICAL FORM

Powder for injection

Cipla Hydrocortisone is white to off-white lyophilized cake/powder in clear tubular glass vial with lyo rubber stopper and yellow coloured flip-off aluminium seal.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

When oral therapy is not feasible, and the strength, form and route of administration of the drug reasonably lend the preparation to the treatment of the condition, Cipla Hydrocortisone powder for injection is indicated for intravenous or intramuscular use in the following conditions:

1. Endocrine disorders

- Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; synthetic analogues may be used in conjunction with mineralocorticoids where applicable; in infancy, mineralocorticoid supplementation is of particular importance). Acute adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; mineralocorticoid supplements may be necessary, particularly when synthetic analogues are used).
- Preoperatively and in the event of serious trauma or illness, in patients with known adrenal insufficiency or when adrenocortical reserve is doubtful
- Shock unresponsive to conventional therapy if adrenocortical insufficiency exists or is suspected
- Congenital adrenal hyperplasia
- Nonsuppurative thyroiditis
- Hypercalcaemia associated with cancer.

2. Rheumatic disorders

As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in:

- Post-traumatic osteoarthritis
- Synovitis of osteoarthritis
- Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low dose maintenance therapy)
- Acute and subacute bursitis
- Epicondylitis
- Acute nonspecific tenosynovitis
- Acute gouty arthritis
- Psoriatic arthritis
- Ankylosing spondylitis.

3. Collagen diseases

During an exacerbation or as maintenance therapy in selected cases of:

- Systemic lupus erythematosus
- Systemic dermatomyositis (polymyositis)
- Acute rheumatic carditis.

4. Dermatological diseases

- Pemphigus
- Severe erythema multiforme (Stevens-Johnson Syndrome)
- Exfoliative dermatitis
- Bullous dermatitis herpetiformis
- Severe seborrhoeic dermatitis
- Severe psoriasis
- Mycosis fungoides.

5. Allergic states

Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in:

- Bronchial asthma

- Drug hypersensitivity reactions
- Contact dermatitis
- Urticarial transfusion reactions
- Atopic dermatitis
- Serum sickness
- Acute noninfectious laryngeal oedema (adrenaline is the drug of first choice).

6. Ophthalmic diseases

Severe acute and chronic allergic and inflammatory processes involving the eye, such as:

- Herpes zoster ophthalmicus
- Iritis, iridocyclitis
- Chorioretinitis
- Diffuse posterior uveitis and choroiditis
- Optic neuritis
- Sympathetic ophthalmia
- Anterior segment inflammation
- Allergic conjunctivitis
- Allergic corneal marginal ulcers
- Keratitis.

7. Gastrointestinal diseases

To tide the patient over a critical period of the disease in:

- Ulcerative colitis (systemic therapy)
- Regional enteritis (systemic therapy).

8. Respiratory diseases

- Symptomatic sarcoidosis
- Loeffler's Syndrome not manageable by other means
- Berylliosis
- Fulminating or disseminated pulmonary tuberculosis when used concurrently with appropriate antituberculous chemotherapy
- Aspiration pneumonitis.

9. Haematological disorders

- Acquired (autoimmune) haemolytic anaemia
- Erythroblastopenia (RBC anaemia)
- Idiopathic thrombocytopenic purpura in adults (IV only; IM administration is contraindicated)
- Secondary thrombocytopenia in adults
- Congenital (erythroid) hypoplastic anaemia.

10. Neoplastic diseases

For palliative management of:

- Leukaemias and lymphomas in adults
- Acute leukaemia in childhood.

11. Oedematous states

- To induce diuresis or remission of proteinuria in the nephrotic syndrome, without uraemia, of the idiopathic type or that due to lupus erythematosus.

12. Miscellaneous

- Tuberculous meningitis with subarachnoid block or impending block when used concurrently with appropriate antituberculous chemotherapy
- Trichinosis with neurological or myocardial involvement.

4.2 Dose and method of administration

Posology

Infants

Formulations containing benzyl alcohol are contraindicated for use in premature infants (see section 4.3 Contraindications).

Adults

This preparation may be administered by intravenous injection or infusion, or by intramuscular injection. The preferred method for initial emergency use is intravenous injection. Following the initial emergency period, consideration should be given to employing a longer acting injectable preparation or an oral preparation.

Therapy is initiated by administering Cipla Hydrocortisone powder for injection intravenously over a period of 30 seconds (e.g., 100 mg) to 10 minutes (e.g., 500 mg or more).

Dosage requirements are variable and must be individualised on the basis of the disease under treatment, its severity and the response of the patient over the entire duration of treatment. A risk/benefit decision must be made in each individual case on an ongoing basis.

The lowest possible dose of corticosteroid should be used to control the condition under treatment for the minimum period. The proper maintenance dosage should be determined by decreasing the initial

drug dosage in small decrements at appropriate time intervals until the lowest dosage, which will maintain an adequate clinical response, is reached.

In general, high dose corticosteroid therapy should be continued only until the patient's condition has stabilised - usually not beyond 48 to 72 hours. Although adverse effects associated with high dose, short-term corticoid therapy are uncommon, peptic ulceration may occur. Prophylactic antacid therapy may be indicated.

When high dose hydrocortisone therapy must be continued beyond 48 to 72 hours, hypernatraemia may occur. Under such circumstances it may be desirable to replace hydrocortisone sodium succinate with a corticoid such as methylprednisolone sodium succinate which causes little or no sodium retention.

If after long-term therapy the drug is to be stopped, it needs to be withdrawn gradually rather than abruptly (see section 4.4 Special warnings and precautions for use).

The initial dose of hydrocortisone sodium succinate powder for injection is 100 mg to 500 mg or more, depending on the severity of the condition. This dose may be repeated at intervals of 2, 4 or 6 hours as indicated by the patient's response and clinical condition. While the dose may be reduced for infants and children, it is governed more by the severity of the condition and response of the patient than by age or body weight but should not be less than 25 mg daily.

Patients subjected to severe stress following corticosteroid therapy should be closely observed for signs and symptoms of adrenocortical insufficiency.

Corticoid therapy is an adjunct to, and not a replacement for, conventional therapy.

Dosage adjustment in hepatic impairment

In patients with liver disease, there may be an increased effect of hydrocortisone resulting from decreased metabolism and elimination of the drug, and reduced dosing may be considered. Monitoring the clinical response to hydrocortisone in these patients should be considered (see section 4.4 Special warnings and precautions for use).

Method of administration

Preparation of solutions

For intravenous or intramuscular injection

Prepare solution by aseptically adding **not more than 2 mL** of Bacteriostatic Water for Injections or Bacteriostatic Sodium Chloride Injection to the contents of one vial.

For intravenous infusion

First prepare solution by adding **not more than 2 mL** of Bacteriostatic Water for Injections to the vial; this solution may then be added to 100 to 1000 mL of the following: 5% glucose in water (or isotonic saline solution or 5% glucose in isotonic saline solution if patient is not on sodium restriction).

Use in one patient on one occasion only. The 100 mg plain vial does not contain an antimicrobial agent. Use solution immediately and discard any residue.

Use diluted/reconstituted solution as soon as possible to avoid microbial contamination hazards, and only if it is clear. If storage is necessary, unused solution may be stored at 2°-8°C for not more than 24 hours provided aseptic procedures are followed. Any solution not used within 24 hours should be discarded.

4.3 Contraindications

Hydrocortisone sodium succinate is contraindicated:

- in patients who have systemic fungal infections.
- in patients with known hypersensitivity to the drug or any component of the formulation.
- for use by the intrathecal route of administration.
- for use by the epidural route of administration.

Administration of live or live attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of corticosteroids (also see section 4.4 Special warnings and precautions for use).

Some Water for Injection may contain benzyl alcohol as a bacteriostatic agent. Benzyl alcohol has been reported to be associated with a fatal “Gaspings Syndrome” in premature infants. The hydrocortisone sodium succinate 100 mg plain vial **DO NOT** contain benzyl alcohol.

Hydrocortisone sodium succinate is not indicated for intrathecal, epidural or local injection, or any other unspecified route of administration.

4.4 Special warnings and precautions for use

Immunosuppressant effects/increased susceptibility to infections

Corticosteroids may increase susceptibility to infection, may mask some signs of infection, and new infections may appear during their use. There may be decreased resistance and inability to localise infection when corticosteroids are used. Infections with any pathogen including viral, bacterial, fungal, protozoan or helminthic infections, in any location in the body, may be associated with the use of corticosteroids alone or in combination with other immunosuppressive agents that affect cellular immunity, humoral immunity, or neutrophil function. These infections may be mild but can be severe and at times fatal. With increasing doses of corticosteroids, the rate of occurrence of infectious complications increases.

Persons who are on drugs which suppress the immune system are more susceptible to infections than healthy individuals. Chicken pox and measles, for example, can have a more serious or even fatal course in non-immune children or adults on corticosteroids.

Administration of live or live attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of corticosteroids. Killed or inactivated vaccines may be administered to patients receiving immunosuppressive doses of corticosteroids; however, the response to such vaccines may be diminished. Indicated immunisation procedures may be undertaken in patients receiving non-immunosuppressive doses of corticosteroids.

The use of hydrocortisone sodium succinate in active tuberculosis should be restricted to those cases of fulminating or disseminated tuberculosis in which the corticosteroid is used for the management of the disease in conjunction with appropriate antituberculosis regimen. If corticosteroids are indicated in patients with latent tuberculosis or tuberculin reactivity, close observation is necessary as reactivation of the disease may occur. During prolonged corticosteroid therapy, these patients should receive chemoprophylaxis.

Kaposi's sarcoma has been reported to occur in patients receiving corticosteroid therapy. Discontinuation of corticosteroids may result in clinical remission.

Controlled clinical trials have failed to establish the efficacy of methylprednisolone sodium succinate in the treatment of sepsis syndrome and septic shock. Two studies suggest that treatment of these conditions with methylprednisolone sodium succinate may increase the risk of mortality in certain patients (i.e., patients with elevated serum creatinine levels or patients who develop secondary infections after receiving methylprednisolone sodium succinate). Although this trial used methylprednisolone sodium succinate only, trial recommends that hydrocortisone sodium succinate not be used for septic shock or sepsis syndrome either.

Immune system effects

Allergic reactions may occur. Because rare instances of skin reactions and anaphylactic/anaphylactoid reactions (e.g., bronchospasm) have occurred in patients receiving parenteral corticosteroid therapy, appropriate precautionary measures should be taken prior to administration, especially when the patient has a history of allergy to any drug.

Endocrine effects

In patients on corticosteroid therapy subjected to unusual stress, increased dosage of rapidly acting corticosteroids before, during and after the stressful situation is indicated.

Pharmacologic doses of corticosteroids administered for prolonged periods may result in hypothalamic-pituitary-adrenal (HPA) suppression (secondary adrenocortical insufficiency). The degree and duration of adrenocortical insufficiency produced is variable among patients and depends on the dose, frequency, time of administration, and duration of corticosteroid therapy.

In addition, acute adrenal insufficiency leading to a fatal outcome may occur if corticosteroids are withdrawn abruptly.

Drug-induced secondary adrenocortical insufficiency may therefore be minimised by gradual reduction of dosage. This type of relative insufficiency may persist for months after discontinuation of therapy, therefore, in any situation of stress occurring during that period, hormone therapy should be reinstated. Since mineralocorticoid secretion may be impaired, salt and/or a mineralocorticoid should be administered concurrently.

A steroid “withdrawal syndrome”, seemingly unrelated to adrenocortical insufficiency, may also occur following abrupt discontinuance of corticosteroids. This syndrome includes symptoms such as: anorexia, nausea, vomiting, lethargy, headache, fever, joint pain, desquamation, myalgia, weight loss, and/or hypotension. These effects are thought to be due to the sudden change in glucocorticoid concentration rather than to low corticosteroid levels.

High doses of corticosteroids can produce or aggravate Cushing’s syndrome. Careful consideration and/or consultation with an endocrinologist are recommended when administering hydrocortisone to patients with Cushing’s disease.

There is an enhanced effect of corticosteroids in patients with hypothyroidism.

Metabolism and nutrition

Corticosteroids, including hydrocortisone, can increase blood glucose, worsen pre-existing diabetes, and predispose those on long-term corticosteroid therapy to diabetes mellitus.

Psychiatric effects

Psychic derangements may appear when corticosteroids are used, ranging from euphoria, insomnia, mood swings, personality changes and severe depression to frank psychotic manifestations. Also, existing emotional instability or psychotic tendencies may be aggravated by corticosteroids.

Potentially severe psychiatric adverse reactions may occur with systemic steroids. Symptoms typically emerge within a few days or weeks of starting treatment. Most reactions recover after either dose reduction or withdrawal, although specific treatment may be necessary. Psychological effects have been reported upon withdrawal of corticosteroids; the frequency is unknown. Patients/caregivers should be encouraged to seek medical attention if psychological symptoms develop in the patient, especially if depressed mood or suicidal ideation is suspected. Patients/caregivers should be alert to possible psychiatric disturbances that may occur either during or immediately after dose tapering/withdrawal of systemic steroids.

Nervous system effects

Corticosteroids should be used with caution in patients with seizure disorders.

Corticosteroids should be used with caution in patients with myasthenia gravis (also see myopathy statement in Musculoskeletal effects).

Severe medical events have been reported in association with the intrathecal/epidural routes of administration.

There have been reports of epidural lipomatosis in patients taking corticosteroids, typically with long-term use at high doses.

Ocular effects

Corticosteroids should be used cautiously in patients with ocular herpes simplex because of possible risk of corneal perforation.

Prolonged use of corticosteroids may produce posterior subcapsular cataracts and nuclear cataracts (particularly in children), exophthalmos, or increased intraocular pressure, which may result in glaucoma with possible damage to the optic nerves. Establishment of secondary fungal and viral infections of the eye may also be enhanced in patients receiving corticosteroids.

Corticosteroid therapy has been associated with central serous chorioretinopathy, which may lead to retinal detachment.

If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes.

Cardiac effects

Adverse effects of glucocorticoids on the cardiovascular system, such as dyslipidaemia and hypertension, may predispose treated patients with existing cardiovascular risk factors to additional cardiovascular effects, if high doses and prolonged courses are used. Accordingly, corticosteroids should be employed judiciously in such patients and attention should be paid to risk modification and additional cardiac monitoring if needed. Low dose therapy may reduce the incidence of complications in corticosteroid therapy.

Systemic corticosteroids should be used with caution, and only if strictly necessary, in patients with congestive heart failure.

Hypertrophic cardiomyopathy has been reported after administration of hydrocortisone to prematurely born infants, therefore appropriate investigation should be undertaken in order to monitor cardiac function and structure in this patient population.

Vascular effects

Thrombosis including venous thromboembolism has been reported to occur with corticosteroids. As a result, corticosteroids should be used with caution in patients who have or may be predisposed to thromboembolic disorders.

Steroids should be used with caution in patients with hypertension.

Gastrointestinal effects

High doses of corticosteroids may produce acute pancreatitis.

There is no universal agreement on whether corticosteroids *per se* are responsible for peptic ulcers encountered during therapy; however, corticosteroid therapy may mask the symptoms of peptic ulcer so that perforation or haemorrhage may occur without significant pain. Glucocorticoid therapy may mask peritonitis or other signs or symptoms associated with gastrointestinal disorders such as perforation, obstruction or pancreatitis. In combination with nonsteroidal anti-inflammatory drugs (NSAIDs), the risk of developing gastrointestinal ulcers is increased.

Corticosteroids should be used with caution in nonspecific ulcerative colitis, if there is a probability of impending perforation, abscess or other pyogenic infection, diverticulitis, fresh intestinal anastomoses, or active or latent peptic ulcer.

Musculoskeletal effects

An acute myopathy has been described with the use of high doses of corticosteroids, most often occurring in patients with disorders of neuromuscular transmission (e.g., myasthenia gravis), or in patients receiving concomitant therapy with anticholinergics, such as neuromuscular blocking drugs (e.g., pancuronium). This acute myopathy is generalised, may involve ocular and respiratory muscles, and may result in quadriplegia. Elevations of creatine kinase may occur. Clinical improvement or recovery after stopping corticosteroids may require weeks to years.

Corticosteroids should be used with caution in patients with osteoporosis.

Investigations

Hydrocortisone can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium. Dietary salt restriction and potassium supplementation may be necessary. All corticosteroids increase calcium excretion.

Injury, poisoning and procedural complications

Systemic corticosteroids are not indicated for and should therefore not be used to treat traumatic brain injury. A large multicentre randomised study in patients administered corticosteroid therapy after significant head injury revealed an increased risk of mortality in the corticosteroid group compared to the placebo group.

Other

Since complications of treatment with glucocorticoids are dependent on the size of the dose and the duration of treatment, a risk/benefit decision must be made in each individual case as to dose and duration of treatment as to whether daily or intermittent therapy should be used.

The lowest possible dose of corticosteroids should be used to control the condition under treatment and when reduction in dosage is possible, the reduction should be gradual.

Aspirin and nonsteroidal anti-inflammatory agents should be used cautiously in conjunction with corticosteroids (see section 4.5 Interactions with other medicines and other forms of interactions).

Pheochromocytoma crisis, which can be fatal, has been reported after administration of systemic corticosteroids. Corticosteroids should only be administered to patients with suspected or identified pheochromocytoma after an appropriate risk/benefit evaluation.

Use in hepatic impairment

Hepatobiliary disorders have been reported which may be reversible after discontinuation of therapy. Therefore, appropriate monitoring is required.

There is an enhanced effect of corticosteroids in patients with cirrhosis.

Hydrocortisone may have an increased effect in patients with liver disease since the metabolism and elimination of hydrocortisone is significantly decreased in these patients.

Use in renal impairment

Corticosteroids should be used with caution in patients with renal insufficiency.

Use in the elderly

No data available.

Paediatric use

Growth and development of infants and children on prolonged corticosteroid therapy should be carefully observed.

Growth may be suppressed in children receiving long-term, daily-divided dose of glucocorticoid therapy and use of such a regimen should be restricted to the most serious indications. Alternate-day glucocorticoid therapy usually avoids or minimises this side effect.

Infants and children on prolonged corticosteroid therapy are at special risk from raised intracranial pressure.

High doses of corticosteroids may produce pancreatitis in children.

Effects on laboratory tests

No data available.

4.5 Interactions with other medicines and other forms of interactions

Hydrocortisone is metabolised by 11 β -hydroxysteroid dehydrogenase type 2 (11 β -HSD2) and the cytochrome P450 (CYP) 3A4 enzyme. The CYP3A4 enzyme catalyses 6 β -hydroxylation of steroids, the essential Phase I metabolic step for both endogenous and synthetic corticosteroids. Many other compounds are also substrates of CYP3A4, some of which have been shown to alter glucocorticoid metabolism by induction (upregulation) or inhibition of the CYP3A4 enzyme.

CYP3A4 INHIBITORS - May decrease hepatic clearance and increase the plasma concentrations of hydrocortisone. In the presence of a CYP3A4 inhibitor (e.g., ketoconazole, itraconazole, clarithromycin, and grapefruit juice), the dose of hydrocortisone may need to be decreased to avoid steroid toxicity.

CYP3A4 INDUCERS - May increase hepatic clearance and decrease the plasma concentrations of hydrocortisone. In the presence of a CYP3A4 inducer (e.g., rifampin, carbamazepine, phenobarbital, and phenytoin), the dose of hydrocortisone may need to be increased to achieve the desired response.

CYP3A4 SUBSTRATES - In the presence of another CYP3A4 substrate, the hepatic clearance of hydrocortisone may be affected, with corresponding dosage adjustments required. It is possible that adverse events associated with the use of either drug alone may be more likely to occur with co-administration.

NON-CYP3A4-MEDIATED EFFECTS - Other interactions and effects that occur with hydrocortisone are described in Table 1 below.

Table 1 provides a list and descriptions of the most common and/or clinically important drug interactions or effects with hydrocortisone.

Table 1. Important drug or substance interactions/effects with hydrocortisone

Drug Class or Type – DRUG or SUBSTANCE	Interaction/Effect
Antibacterial – ISONIAZID	CYP3A4 INHIBITOR
Antibiotic, Antitubercular – RIFAMPIN	CYP3A4 INDUCER
Anticoagulants (oral)	The effect of corticosteroids on oral anticoagulants is variable. There are reports of enhanced as well as diminished effects of anticoagulants when given concurrently with corticosteroids. Therefore, coagulation indices should be monitored to maintain the desired anticoagulant effects.
Anticonvulsants – CARBAMAZEPINE	CYP3A4 INDUCER (and SUBSTRATE)
Anticonvulsants – PHENOBARBITAL (PHENOBARBITONE) – PHENYTOIN	CYP3A4 INDUCERS
Anticholinergics – NEUROMUSCULAR BLOCKERS	Corticosteroids may influence the effect of anticholinergics. 1) An acute myopathy has been reported with the concomitant use of high doses of corticosteroids and anticholinergics, such as neuromuscular blocking drugs (see section 4.4 Special warnings and precautions for use - Musculoskeletal effects, for additional information). 2) Antagonism of the neuromuscular blocking effects of pancuronium and vecuronium has been reported in patients taking corticosteroids. This interaction may be expected with all competitive neuromuscular blockers.
Anticholinesterases	Steroids may reduce the effects of anticholinesterases in myasthenia gravis.
Antidiabetics	Because corticosteroids may increase blood glucose concentrations, dosage adjustments of antidiabetic agents may be required.
Antiemetic – APREPITANT – FOSAPREPITANT	CYP3A4 INHIBITORS (and SUBSTRATES)
Antifungals – ITRACONAZOLE – KETOCONAZOLE	CYP3A4 INHIBITORS (and SUBSTRATES)
Antivirals	CYP3A4 INHIBITORS (and SUBSTRATES)

– HIV-PROTEASE INHIBITORS	<ol style="list-style-type: none"> 1) Protease inhibitors, such as indinavir and ritonavir, may increase plasma concentrations of corticosteroids. 2) Corticosteroids may induce the metabolism of HIV-protease inhibitors resulting in reduced plasma concentrations.
Aromatase Inhibitors – AMINOGLUTETHIMIDE	Aminoglutethimide-induced adrenal suppression may exacerbate endocrine changes caused by prolonged glucocorticoid treatment.
Calcium Channel Blocker – DILTIAZEM	CYP3A4 INHIBITOR (and SUBSTRATE)
Cardiac Glycosides – DIGOXIN	Concurrent use of corticosteroids with cardiac glycosides may enhance the possibility of arrhythmias or digitalis toxicity associated with hypokalaemia. In all patients taking any of these drug therapy combinations, serum electrolyte determinations, particularly potassium levels, should be monitored closely.
Contraceptives (oral) – ETHINYLESTRADIOL/ NORETHISTERONE	CYP3A4 INHIBITOR (and SUBSTRATE)
– GRAPEFRUIT JUICE	CYP3A4 INHIBITOR
Immunosuppressant – CICLOSPORIN	CYP3A4 INHIBITOR (and SUBSTRATE) Increased activity of both ciclosporin and corticosteroids may occur when the two are used concurrently. Convulsions have been reported with this concurrent use.
Immunosuppressant – CYCLOPHOSPHAMIDE – TACROLIMUS	CYP3A4 SUBSTRATES
Macrolide Antibacterial – CLARITHROMYCIN – ERYTHROMYCIN	CYP3A4 INHIBITORS (and SUBSTRATES)
Macrolide Antibacterial – TROLEANDOMYCIN	CYP3A4 INHIBITOR
NSAIDs – high dose ASPIRIN (acetylsalicylic acid)	<ol style="list-style-type: none"> 1) There may be increased incidence of gastrointestinal bleeding and ulceration when corticosteroids are given with NSAIDs. 2) Corticosteroids may increase the clearance of high dose aspirin, which can lead to decreased salicylate serum levels. Discontinuation of corticosteroid treatment can lead to raised salicylate serum levels, which could lead to an increased risk of salicylate toxicity.
Potassium Depleting Agents	When corticosteroids are administered concomitantly with potassium depleting agents (i.e., diuretics), patients should be observed closely for development of hypokalaemia. There is also an increased risk of hypokalaemia with concurrent use of corticosteroids with amphotericin B, xanthines, or beta2 agonists. There have been cases reported in which concomitant use of amphotericin B and hydrocortisone was followed by cardiac enlargement and congestive heart failure.

The pharmacokinetic interactions listed below are potentially clinically important.

1. Oral contraceptives retard the metabolism of hydrocortisone due to its increased binding to globulin (transcortin). This increases the plasma levels of hydrocortisone thus potentiating its biological effect. Dosage adjustments of hydrocortisone may be required if estrogens are added to or withdrawn from a stable dosage regimen.

2. Drugs that induce hepatic enzymes such as phenobarbitone, phenytoin and rifampicin may increase the clearance of corticosteroids and may require increases in corticosteroid dose to achieve the desired response.
3. Drugs such as troleandomycin and ketoconazole may inhibit the metabolism of corticosteroids and thus decrease their clearance. Therefore, the dose of corticosteroid should be titrated to avoid steroid toxicity.

4.6 Fertility, pregnancy and lactation

Effects on fertility

No specific animal or clinical studies on the effects of hydrocortisone on fertility have been performed. Corticosteroids have been shown to impair fertility and reduce embryonic viability in studies in mice and rats.

Use in pregnancy – Pregnancy Category C

Since adequate human reproductive studies have not been done with hydrocortisone sodium succinate, this medicinal product should be used during pregnancy only after a careful assessment of the benefit-risk ratio to the mother and fetus.

In animal experiments, corticosteroids have been found to cause malformations of various kinds (cleft palate, skeletal malformations) and abortion. These findings do not seem to be relevant to human beings. Some corticosteroids readily cross the placenta. Reduced placental and birth weight have been recorded in animals and humans after long-term treatment. Since the possibility of suppression of the adrenal cortex in the newborn baby after long-term treatment must be considered, the needs of the mother must be carefully weighed against the risk to the fetus when prescribing corticosteroids. Some retrospective studies have found an increased incidence of low birth weights in infants born of mothers receiving corticosteroids. In humans, the risk of low birth weight appears to be dose-related and may be minimised by administering lower corticosteroid doses. The short-term use of corticosteroids antepartum for the prevention of respiratory distress syndrome does not seem to pose a risk to the fetus or the newborn infant.

Infants born of mothers who have received substantial doses of corticosteroids during pregnancy must be carefully observed and evaluated for signs of adrenal insufficiency.

Maternal pulmonary oedema has been reported with tocolysis and fluid overload.

Cataracts have been observed in infants born to mothers treated with long-term corticosteroids during pregnancy.

Use in lactation

Prednisolone is excreted in breast milk, therefore it is reasonable to assume that all corticosteroids are. No specific data are known for hydrocortisone sodium succinate. Therefore, it is recommended that breastfeeding should cease in women who will be or are receiving corticosteroids.

4.7 Effects on ability to drive and use machines

The effect of corticosteroids on the ability to drive or use machinery has not been systematically evaluated. Undesirable effects, such as syncope, vertigo and convulsions are possible after treatment with corticosteroids (see section 4.8 Adverse effects (undesirable effects)). If affected, patients should not drive or operate machinery.

4.8 Adverse effects (undesirable effects)

Infections and infestations

- Infection masked opportunistic infections (with any pathogen, in any location in the body, from mild to fatal)
- Infections (becoming active, including reactivation of tuberculosis).

Neoplasms benign, malignant and unspecified (including cysts and polyps)

- Kaposi's sarcoma (has been reported to occur in patients receiving corticosteroid therapy).

Blood and lymphatic system disorders

- Leucocytosis.

Immune system disorders

- Drug hypersensitivity
- Anaphylactic reaction
- Anaphylactoid reaction.

Endocrine disorders

- Cushingoid
- Hypopituitarism
- Steroid withdrawal syndrome
- Manifestations of latent diabetes mellitus.

Metabolism and nutrition disorders

- Metabolic acidosis
- Sodium retention
- Fluid retention
- Alkalosis hypokalaemic
- Dyslipidaemia
- Glucose tolerance impaired
- Increased insulin requirement (or oral hypoglycaemic agents in diabetics)
- Lipomatosis
- Increased appetite (which may result in weight increased).

Psychiatric disorders

- Affective disorder (including depression, euphoric mood, affect lability, drug dependence, suicidal ideation)
- Psychotic disorder (including mania, delusion, hallucination and schizophrenia)
- Mental disorder
- Personality change
- Confusional state
- Anxiety
- Mood swings
- Abnormal behaviour
- Insomnia
- Irritability.

Nervous system disorders

- Epidural lipomatosis
- Intracranial pressure increased
- Benign intracranial hypertension
- Seizure
- Amnesia
- Cognitive disorder
- Dizziness
- Headache.

Eye disorders

- Central serous chorioretinopathy
- Cataract
- Glaucoma
- Exophthalmos
- Vision blurred.

Ear and labyrinth disorders

- Vertigo.

Cardiac disorders

- Congestive heart failure (in susceptible patients)
- Hypertrophic cardiomyopathy in prematurely born infants.

Vascular disorders

- Thrombosis
- Hypertension
- Hypotension.

Respiratory, thoracic and mediastinal disorders

- Pulmonary embolism
- Gasping Syndrome
- Hiccups.

Gastrointestinal disorders

- Peptic ulcer (with possible peptic ulcer perforation and peptic ulcer haemorrhage)
- Intestinal perforation
- Gastric haemorrhage
- Pancreatitis
- Oesophagitis
- Abdominal distension
- Abdominal pain
- Diarrhoea
- Dyspepsia
- Nausea.

Skin and subcutaneous tissue disorders

- Angioedema
- Hirsutism
- Petechiae
- Ecchymosis
- Skin atrophy
- Erythema

- Hyperhidrosis
- Skin striae
- Rash
- Pruritus
- Urticaria
- Acne
- Skin hypopigmentation.

Musculoskeletal and connective tissue disorders

- Muscle weakness
- Myalgia
- Myopathy
- Muscle atrophy
- Osteoporosis
- Osteonecrosis
- Pathological fracture
- Neuropathic arthropathy
- Arthralgia
- Growth retardation.

Reproductive system and breast disorders

- Menstruation irregular.

General disorders and administration site conditions

- Impaired healing
- Oedema peripheral
- Fatigue
- Malaise
- Injection site reaction.

Investigations

- Intraocular pressure increased

- Carbohydrate tolerance decreased
- Blood potassium decreased
- Urine calcium increased
- Alanine aminotransferase increased, aspartate aminotransferase increased, blood alkaline phosphatase increased. These changes are usually small, not associated with any clinical syndrome and are reversible upon discontinuation.
- Blood urea increased
- Suppression of reactions to skin tests.

Injury, poisoning and procedural complications

- Spinal compression fracture
- Tendon rupture.

The following additional reactions are related to parenteral corticosteroid therapy:

- Hyperpigmentation or hypopigmentation
- Subcutaneous and cutaneous atrophy
- Sterile abscess.

Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at <https://www.tga.gov.au/reporting-problems>.

4.9 Overdose

Symptoms and signs

Reports of acute toxicity and metabolic disturbances with glucocorticoids are rare but do occur. There is no clinical syndrome of acute overdosage with hydrocortisone sodium succinate. Acute overdose may possibly aggravate pre-existing disease states such as ulceration of the gastrointestinal tract, electrolyte disturbances, infections, diabetes and oedema.

Repeated frequent doses (daily or several times per week) over a protracted period may result in a Cushingoid state. The possibility of adrenal suppression should be guarded against by gradual diminution of dose levels over a period of time.

Treatment

In the event of acute overdose, treatment is symptomatic and supportive, including respiratory and cardiovascular function. In chronic toxicity, fluids and electrolytes should be monitored closely. Serum levels are not clinically useful.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Mechanism of action

Hydrocortisone sodium succinate is an anti-inflammatory adrenocortical steroid. This highly water-soluble sodium succinate ester of hydrocortisone permits the immediate intravenous administration of high doses of hydrocortisone in a small volume of diluent and is particularly useful where high blood levels of hydrocortisone are required rapidly.

Hydrocortisone sodium succinate has the same metabolic and anti-inflammatory actions as hydrocortisone. When given parenterally and in equimolar quantities, the two compounds are equivalent in biological activity.

Clinical trials

No data available.

5.2 Pharmacokinetic properties

Absorption

Following the intravenous injection of hydrocortisone sodium succinate, demonstrable effects are evident within 1 hour and persist for a variable period. This preparation is also rapidly absorbed when administered intramuscularly. Thus, if constantly high blood levels are required, injections should be made every 4 to 6 hours.

Metabolism

Hydrocortisone is metabolised by 11 β -hydroxysteroid dehydrogenase type 2 (11 β -HSD2) and the cytochrome P450 (CYP) 3A4 enzyme. The CYP3A4 enzyme catalyses 6 β -hydroxylation of steroids, the essential Phase I metabolic step for both endogenous and synthetic corticosteroids.

Excretion

Excretion of the intravenously administered dose is nearly complete within 12 hours. Intramuscular injections are excreted in a pattern similar to that observed after intravenous injections.

5.3 Preclinical safety data

Genotoxicity

No data available.

Carcinogenicity

No data available.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Dibasic sodium phosphate,
Monobasic sodium phosphate,

Sodium hydroxide (for pH adjustment),
Water for injection.

6.2 Incompatibilities

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

6.3 Shelf life

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

6.4 Special precautions for storage

Store unconstituted powder below 25°C; protect from light.

6.5 Nature and contents of container

Cipla Hydrocortisone is filled in 5 mL USP Type-I Fiolax Clear Tubular EBB lyophilised vial with 20 mm grey bromobutyl stopper and 20 mm yellow Al flip-off seal.

Cipla Hydrocortisone is available in the following pack size:

5 x 100 mg powder for injection in glass vials.


6.6 Special precautions for storage

In Australia, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

6.7 Physicochemical properties

Description

Hydrocortisone sodium succinate is a white to almost white, hygroscopic powder. It is practically insoluble in water, freely soluble in acetone and in anhydrous alcohol (ethanol).

Chemical structure:	
Non-proprietary name:	Hydrocortisone sodium succinate
Chemical name:	pregn-4-ene-3,20-dione,21-(3-carboxy-1-oxopropoxy)-11,17-dihydroxy-, monosodium salt, (11β)-
Molecular formula:	C ₂₅ H ₃₃ O ₈ Na
Molecular weight:	484.52

CAS number

125-04-2

7. MEDICINE SCHEDULE (POISONS STANDARD)

Schedule 4, Prescription Only Medicine

8. SPONSOR

Cipla Australia Pty Ltd.,
Level 1, 132-136 Albert Road,
South Melbourne, VIC 3205.
drugsafety@cipla.com
Toll Free number: 1800 569 074

9. DATE OF FIRST APPROVAL

08 June 2022

10. DATE OF REVISION

Not Applicable

Summary table of changes

Section changed	Summary of new information
NA	NA

NA – Not Applicable